


Volume 13, 18 December 2013

Publisher: Igitur publishing

URL: <http://www.ijic.org>

Cite this as: Int J Integr Care 2013; WCIC Conf Suppl; [URN:NBN:NL:UI:10-1-115971](https://nbn-resolving.org/urn:nbn:nl:ui:10-1-115971)

Copyright: 

Poster Abstract

Managing mental illness in primary care - General Practitioners' perspective

Wayne Freeman Chong, Senior Research Analyst, Agency for Integrated Care, Singapore

Harold Lim, Senior Executive, Agency for Integrated Care, Singapore

Shing Yew Joel Yong, Senior Executive, Agency for Integrated Care, Singapore

Correspondence to: **Mr Wayne Freeman Chong**, Senior Research Analyst, Knowledge Management Division, Agency for Integrated Care, Singapore, E-mail: waynefreeman.chong@aic.sg

Abstract

Background: The Agency for Integrated Care (AIC) conducted a survey to determine the specific areas in which the agency could work with General Practitioners (GPs) to enhance mental illness management in Singapore. This survey was conducted as part of AIC's engagement strategy and because the Primary Care Surveys 2010 and 2005 noted that a sizeable number of mental health cases are seen by GPs.

Methodology: An interviewer-administered questionnaire was conducted among a random sample of 849 group and non-group GPs practicing in four geographical districts of Singapore. The response rate was 64%.

Findings: About 70% of the respondent GPs indicated that they were seeing patients with mental illness. Anxiety and depression were the most commonly seen mental conditions. The lack of available drugs and facilities in the clinic, and unfamiliarity with mental health patient management were the main reasons that GPs do not see patients with mental illness. About 39% of the GPs referred patients to a psychologist. About 36% of the GPs referred patients to a counsellor. Availability of additional allied health services in the neighbourhood was crucial in encouraging GPs to see more patients with mental illness. General geriatrics, dementia, general mental health and major depression were the top four areas of interest for further training.

Conclusions: The role of GPs in managing mental illnesses remains crucial. There is more pressing need to address their perceived difficulties in managing such patients. The results of the GP Landscape Survey provided evidential bases to enhance the National Mental Health Blueprint in three areas: (i) extension of GP consultation subsidies to middle-income patients, (ii) expansion of allied mental health services to support GPs, and (iii) development of training programmes for GPs in mental illnesses that are commonly encountered by and of interest to GPs.

Keywords:

general practice, gp, mental illness, primary care

References:

1. Lum AW, Kwok KW, Chong SA., et al. Providing integrated mental health services in the Singapore primary care setting--the general practitioner psychiatric programme experience. *Ann Acad Med Singapore*. 2008;37:128-31.
2. Sng QS. Singapore Ministry of Health. Primary Care Survey 2010. Profile of Primary Care Patients. 2011. Assessed Nov 26 2012. Available at http://www.moh.gov.sg/content/dam/moh_web/Publications/Information%20Papers/2011/Primary%20Care%20Survey%202010%20-%20Profile%20of%20Primary%20Care%20Patients.pdf.
3. Singapore Ministry of Health. Primary Care Survey 2005. 2006. Accessed Nov 26 2012. Available at http://www.moh.gov.sg/content/dam/moh_web/Publications/Reports/2006/1/Primary_Care_Survey_2005_0.pdf.
4. Chong SA, Abdin E, Vaingankar JA, et al. A Population-based Survey of Mental Disorders in Singapore. *Ann Acad Med*. 2012;41(2):49-66.
5. US Department of Health and Human Services. Mental Health: A Report of the Surgeon General. Rockville, MD: US Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health, 1999; p. 13-20.
6. Goldman HH, Adams NH, Taube, CA. Deinstitutionalization: the data demythologized. *Hosp Com Psychiatry*. 1983;34:124-9.
7. Lum WMA, Chew J, Lim BL. The Successful Collaboration Between Psychiatrists, A Mental Health Institution and General Practitioners in Primary Care. *Singapore Fam Phys*. 2010; 36(4):19-21.
8. Thornicroft G, Tansella M. WHO Regional Office for Europe's Health Evidence Network. What are the arguments for community-based mental health care? August 2003. Accessed Nov 26 2012. Available at http://www.euro.who.int/_data/assets/pdf_file/0019/74710/E82976.pdf.
9. Janhavi AV, Fong CW, Kwok KW, et al. Managing Patients with Mental Illness in Primary Care: Apprehension and Views of General Practitioners. *Singapore Fam Phys*. 2010;36(4): 22-25.
10. Barlett JE, Kotrlík JW, Higgins CC. Organizational Research: Determining Appropriate Sample Size in Survey Research. *Information Technology, Learning and Performance*. 2001; 19(1): 43-50.
11. Alsnih R, Stopher P. Institute of Transport Studies. The University of Sydney. Working Paper. Estimating Eligibility Rates: A Crucial Component of the Calculation for Response Rates. 2004. Accessed Nov 26 2012. Available at http://ws.econ.usyd.edu.au/itls/wp-archive/itls_wp_04-19.pdf.
12. King M, Nazareth I, Levy G, et al. Prevalence of common mental disorders in general practice attendees across Europe. *Brit J Psychiatry*. 2008;192:362-7.
13. Comparelli A, Lazanio S, Tatarelli R. Psychiatric disorders and their management in general practice. *Psychiatr Serv*. 2001;52:388-9.
14. Singapore Ministry of Health. Committee of Supply Speech for Minister Of State for Health. 28 Mar 2012. Community Health: Working with and through the Community for Better Health Outcomes. Accessed Nov 26 2012. Available at http://www.moh.gov.sg/content/moh_web/home/pressRoom/speeches_d/2012/COS_speech_MOS_Health_-_Community_Health.html.

15. Agency for Integrated Care, Singapore. Temasek Cares – iCommunity@North: First regional mental health network launched in the Northern region of Singapore. 30 Jul 2012. Accessed 26 Nov 2012. Available at <http://www.aic.sg/page.aspx?id=2147484147>
